

Spatial image of reaction area from scattering. II :

On connection between the differential cross-sections in transverse momentum and in nearest approach parameter.

N.Bobrovskaya and A.N.Vall*

Department of Theoretical Physics, Irkutsk State University, Irkutsk, 664003 Russia

M.V.Polyakov[†]

Institut für Theoretische Physik II Ruhr-Universität Bochum, NB6 D-44780 Bochum, Germany

A.A.Vladimirov[‡]

Bogoliubov Laboratory of Theoretical Physics, JINR, 141980, Moscow Region, Dubna, Russia

The connection between differential cross sections of particle C creation on transverse momentum and on nearest approach parameter $\vec{b}$ is investigated in the context of formalism $SO(2,1)$ algebra. Where parameter $\vec{b}$ characterizes particle creation area. This distribution tightly concerned with spatial structure of particles interaction and allows intuitive physic interpretation. It is shown that area of large transverse momentum carries in the main contribution to distribution function on b in backward semisphere in interval $1/2 \leq bq \sim 1$, where q is momentum of the particle C . The left border of inequation defines by Geizenberg uncertainty relation on parameters "momentum - radius of localization area". The spatial structure of creation area in transverse momentum plane is a set of discrete axially-symmetrical (for spinless particles C) zones. The received connection between cross sections on $\vec{q}_\perp$ and $\vec{b}$ is exact and don't connect with any model. So ability appears to analyze the spatial structure of target using experimental data of particle C angle distribution. It was received an exact relation between $\langle b_\perp^2 \rangle$ and $\langle \cos^2 \theta_\pm \rangle$ for any $A + B \rightarrow C + D$ processes in center of particles A and B mass frame, where average is going on corresponding differential cross-sections. It is shown that quantum-mechanic constrain on b^2 spectrum ($b^2 q^2 > \hbar^2/4$) brings to constrain $\langle \cos^2 \theta_\pm \rangle \geq 1/8$. As application it is considered the process $\gamma + p \rightarrow \pi^0 + p$ at photon energy $E_\gamma = 5\text{Gev}$.

PACS numbers:

DIFFERENTIAL CROSS-SECTION ON THE TRANSVERSE MOMENTUM OF PARTICLE C

In this paper we will reproduce in detail the derivation of connection between the differential cross-section in particle C momentum and S -matrix element [1]. As in the framework of the same scheme the similar calculation for the cross-section on nearest approach parameter b [2] will be made.

Let us examine the process $A + B \rightarrow C + D$ for two physical cases. In the first case particle C is created with fixed momentum $\vec{q}$, the second, particle C is created in state $|\vec{\mu}, q, \epsilon\rangle$. The differential cross-section on transverse momentum of the particle C is

$$\frac{d\sigma^\pm}{d\Omega_{\vec{q}}} = \frac{1}{n_1 n_2 T V |\vec{u}|} \frac{dN^\pm}{d\Omega_{\vec{q}}} . \quad (1)$$

Here $\vec{u}$ is relative speed of initial particles, n_1, n_2 are densities of particles A and B , $(\pm)$ means that the particle C scatters to the forward and backward half-sphere (z -axis is directed along the momentum of particle A), and N is the total number of particles C and D creation events in all space (volume V) during infinite time (T). This is

$$\begin{aligned} N &= \int d\vec{q} d\vec{q}_1 | \langle \vec{q}; \vec{q}_1 | \hat{F} | in \rangle |^2 = \sum_{\epsilon=\pm 1} N^{(\epsilon)} = \\ &= \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{q}} | \langle \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 | \hat{F} | in \rangle |^2 , \end{aligned} \quad (2)$$

where

$$|in\rangle = (2\pi)^3 n_1^{1/2} n_2^{1/2} a^+(\vec{p}_1) a^+(\vec{p}_2) |0\rangle \equiv (2\pi)^3 n_1^{1/2} n_2^{1/2} |\vec{p}_1; \vec{p}_2\rangle .$$

The creation operators are related to particles A and B , operator $\hat{F}$ is connected with S -matrix by relation $S = I + i\hat{F}$. Translational invariance allows present the matrix element in the form:

$$\langle f|\hat{F}|in \rangle = \delta^{(4)}(q_{in} - q_f) \langle f|\hat{A}|in \rangle, \quad (3)$$

therefore:

$$\begin{aligned} N &= \frac{TV}{(2\pi)^4} \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{q}} \delta^{(4)}(q + q_1 - P) \times \\ &\times |\langle \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 | \hat{A} | in \rangle|^2 = \\ &= \frac{TV}{(2\pi)^4} \sum_{\epsilon=\pm 1} \int q^2 dq d\Omega_{\vec{q}} \delta(E_q + \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} - P^0) \times \\ &\times |\langle \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 = \vec{P} - \vec{q} | \hat{A} | in \rangle|^2. \end{aligned} \quad (4)$$

Here, the four-momentum $P = P_1 + P_2$. In (4) the δ -function determines the energy shell for particle C reaction:

$$\left\{ E_q + \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} - P^0 \right\}_{q=\tilde{q}} = 0. \quad (5)$$

Let us define:

$$\begin{aligned} E_D &= \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} \\ \lambda(\vec{q}_\perp) &= \left\{ \frac{q E_q E_D}{q^2 P^0 - (\vec{q} \cdot \vec{P}) E_q} \right\}_{q=\tilde{q}}. \end{aligned} \quad (6)$$

Then the following relation is valid:

$$\delta(E_q + \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} - P^0) = \lambda(\vec{q}_\perp) \delta(q - \tilde{q}). \quad (7)$$

Integrating (4) over q we get:

$$\begin{aligned} N &= \frac{TV}{(2\pi)^4} \sum_{\epsilon=\pm 1} \int d\Omega_{\vec{q}} q^2 \lambda(\vec{q}_\perp) \times \\ &\times |\langle \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 = \vec{P} - \vec{q} | A | in \rangle|^2, \quad q = \tilde{q}, \quad q = |\vec{q}|. \end{aligned} \quad (8)$$

So, we obtain a relation for particle C transverse momentum distribution:

$$\frac{dN^\pm}{d\Omega_{\vec{q}}} = \frac{TV}{(2\pi)^4} q^2 \lambda(\vec{q}_\perp) |\langle \vec{q}_\perp, \pm \sqrt{q^2 - q_\perp^2}; \vec{q}_1 = \vec{P} - \vec{q} | A | in \rangle|^2, \quad q = \tilde{q}. \quad (9)$$

Let us define:

$$A^{(\epsilon)}(\vec{q}) = \langle \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 = \vec{P} - \vec{q} | A | \vec{p}_1; \vec{p}_2 \rangle, \quad q = \tilde{q}. \quad (10)$$

Then (9) takes a view:

$$\frac{dN^\pm}{d\Omega_{\vec{q}}} = (2\pi)^2 TV n_1 n_2 q^2 \lambda(\vec{q}_\perp) |A^{(\pm)}(\vec{q})|^2, \quad q = \tilde{q}. \quad (11)$$

The differential cross-section is resulted by substitution (11) to relation (1).

Here we mark an important notice. An angle dependents in cross section $\frac{d\sigma^\pm}{d\Omega_{\vec{q}}}$ appears not only through variable $\vec{q}_\perp$, but in general case also through $\tilde{q}$. In the frame of particles A and B centre of mass (c.m.f.) the particle C energy defined only through square of invariant mass $s = (p_A + p_B)^2$. And it is

$$E_C^* = \frac{1}{2\sqrt{s}}(s + m_C^2 - m_D^2). \quad (12)$$

So:

$$\tilde{q} = p_C^* = \sqrt{(E_C^*)^2 - m_C^2} .$$

And E_C^* isn't depend for scattering angle (the star means c.m.f.). The situation changes in the labor frame (l.f.). In this case there are critical parameter κ which is [3]

$$\kappa = \frac{\sqrt{s} \cdot p_C^*}{m_C \cdot p_A}$$

. In the case $\kappa > 1$ the connection between particle C momentum and scattering angle has a single meaning and has a view:

$$\tilde{q} = p_C = \frac{\sqrt{s} E_C^* p_A \cos(\theta) + (E_A + m_A)[s(p_C^*)^2 - m_C^2 p_A^2 \sin^2(\theta)]^{1/2}}{(E_A + m_B)^2 - p_A^2 \cos^2(\theta)} , \quad (13)$$

where $0 < \theta < \pi$ is scattering angle in l.f. (there isn't the maximum angle limitation). We will back to discuss this expression below.

THE DIFFERENTIAL CROSS-SECTION ON NEAREST APPROACH PARAMETER $\vec{b}$ IN C.M.F.

Now we discuss the case, when the particle C creates in state $|\vec{\mu}, q, \epsilon >$. Choosing for state $|out >$ the corresponding basis, we obtain following relation for the total number of C and D particles creation events [2]:

$$N = (2\pi)^2 \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{\mu}} | < \vec{\mu}, q, \epsilon; \vec{q}_1 | \hat{F} | in > |^2 . \quad (14)$$

As distinct from the previous case, the state $< \vec{\mu}, q, \epsilon; \vec{q}_1 |$ is not a state with a define momentum. So the particle C cross-section in this state can't be expressed through the matrix element $< \vec{\mu}, q, \epsilon; \vec{q}_1 | \hat{F} | in >$. To find $\frac{d\sigma^\pm}{d\Omega_{\vec{\mu}}}$ we will use a state expansion on states with the fixed particle C transverse momentum [2]:

$$< \vec{\mu}, q, \epsilon; \vec{q}_1 | = \frac{1}{(2\pi)^2} \int \bar{\xi}(\vec{q}_\perp, \vec{\mu}) < \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 | d\Omega_{\vec{q}} . \quad (15)$$

Substituting this expansion to (14) we obtain:

$$N = \frac{1}{(2\pi)^2} \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} d\Omega_{\vec{k}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) \times \\ \times F(\vec{q}_\perp, \vec{q}_1) \bar{F}(\vec{k}_\perp, \vec{q}_1) , \quad |\vec{q}| = |\vec{k}| = q , \quad (16)$$

where we use a notation:

$$< \vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}; \vec{q}_1 | \hat{F} | in > = F(\vec{q}_\perp, \vec{q}_1) . \quad (17)$$

Let us make an identical transformation in (16):

$$F(\vec{q}_\perp, \vec{q}_1) = [F(\vec{q}_\perp, \vec{q}_1) - F(\vec{k}_\perp, \vec{q}_1)] + F(\vec{k}_\perp, \vec{q}_1) ,$$

$$\bar{F}(\vec{k}_\perp, \vec{q}_1) = [\bar{F}(\vec{k}_\perp, \vec{q}_1) - \bar{F}(\vec{q}_\perp, \vec{q}_1)] + \bar{F}(\vec{q}_\perp, \vec{q}_1) .$$

Then:

$$N = \frac{1}{(2\pi)^2} Re \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} d\Omega_{\vec{k}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) |F(\vec{q}_\perp, \vec{q}_1)|^2 - \\ - \frac{1}{(2\pi)^2} \frac{1}{2} \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} d\Omega_{\vec{k}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) \times \\ \times |F(\vec{q}_\perp, \vec{q}_1) - F(\vec{k}_\perp, \vec{q}_1)|^2 , \quad |\vec{q}| = |\vec{k}| = q . \quad (18)$$

The second term in (18) is turned to zero, owing to the completeness of the system of basic functions $\xi(\vec{k}_\perp, \vec{\mu})$ [2] :

$$\int d\Omega_{\vec{\mu}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) \sim \delta(\vec{k}_\perp - \vec{q}_\perp) .$$

Finally we obtain:

$$N = \frac{1}{(2\pi)^2} \text{Re} \sum_{\epsilon=\pm 1} \int d\vec{q}_1 q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} d\Omega_{\vec{k}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) |F(\vec{q}_\perp, \vec{q}_1)|^2 . \quad (19)$$

Such representation for N is universal in the meaning that it follows both differential cross-sections in terms of $\vec{q}_\perp$ and also in terms of $\vec{\mu}$. This allows mark out singular factors in N , connected with infinite time and volume. Turning to the matrix elements from $\hat{F}$ to $\hat{A}$ and integrating over momentum $\vec{q}_1$, we get:

$$N = \frac{VT}{(2\pi)^6} \text{Re} \sum_{\epsilon=\pm 1} \int q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} d\Omega_{\vec{k}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \xi(\vec{k}_\perp, \vec{\mu}) \times \\ \times \delta(E_q + \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} - P^0) |A^{(\epsilon)}(\vec{q})|^2 . \quad (20)$$

If we integrate this correlation over the parameter $\vec{\mu}$ and momentum q then we automatically obtain relation (11). But we will integrate over momentum $\vec{k}_\perp$. So we have:

$$N = \frac{VT}{(2\pi)^6} \text{Re} \sum_{\epsilon=\pm 1} \int q^2 dq d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} \kappa(\mu) \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \times \\ \times \delta(E_q + \sqrt{(\vec{q} - \vec{P})^2 + m_D^2} - P^0) |A^{(\epsilon)}(\vec{q})|^2 , \quad (21)$$

where [4]

$$\kappa(\mu) = \int \xi(\vec{k}_\perp, \vec{\mu}) d\Omega_{\vec{k}} = \frac{2\pi^2}{ch(\pi\mu)} \frac{\sqrt{\pi}}{|\Gamma(\frac{i\mu}{2} + \frac{3}{4})|^2} = \sqrt{\pi} \left| \Gamma(\frac{i\mu}{2} + \frac{1}{4}) \right|^2 . \quad (22)$$

Further integrating over q in relation (21) we obtain:

$$N = \frac{VT}{(2\pi)^6} \text{Re} \sum_{\epsilon=\pm 1} \int d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} \kappa(\mu) \bar{\xi}(\vec{q}_\perp, \vec{\mu}) q^2 \lambda(\vec{q}_\perp) |A^{(\epsilon)}(\vec{q})|^2 , q = \tilde{q} . \quad (23)$$

Taking into account (11) we can rewrite this expression in the following form:

$$N = \frac{1}{(2\pi)^2} \text{Re} \sum_{\epsilon=\pm 1} \int d\Omega_{\vec{\mu}} d\Omega_{\vec{q}} \kappa(\mu) \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \frac{dN^{(\epsilon)}}{d\Omega_{\vec{q}}} , q = \tilde{q} . \quad (24)$$

This relation for the total number of events N is right in any frame of particles A and B . Crossing from N to differential distribution demands fixation of frame. If the initial state set in c.m.f, then $\tilde{q}$ don't depend of the scattering angle. And in this case we have for the differential distribution on $\vec{\mu}$:

$$\frac{dN^\pm}{d\Omega_{\vec{\mu}}} = \frac{1}{(2\pi)^2} \kappa(\mu) \text{Re} \int d\Omega_{\vec{q}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \frac{dN^\pm}{d\Omega_{\vec{q}}} , q = \tilde{q} , \quad (25)$$

where

$$\tilde{q}^2 = \frac{(s + m_C^2 - m_D^2)^2}{4s} - m_C^2 .$$

There are $q = \tilde{q}$ everywhere below (reaction surface).

Using relations between μ and b

$$\mu = (b^2 q^2 - \frac{1}{4})^{1/2} , d\Omega_{\vec{\mu}} = q^2 \tanh(\pi\mu) d\vec{b} , d\vec{\mu} = \mu d\mu d\varphi , d\vec{b} = b db d\varphi , \quad (26)$$

we get a distribution on $\vec{b}$. So we have for differential cross-sections:

$$\begin{aligned}\frac{d\sigma^\pm}{d\Omega_{\vec{\mu}}} &= \frac{1}{(2\pi)^2} \kappa(\mu) \operatorname{Re} \int d\Omega_{\vec{q}} \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \frac{d\sigma^\pm}{d\Omega_{\vec{q}}}, \\ \frac{d\sigma^\pm}{d\Omega_{\vec{q}}} &= (2\pi)^2 \frac{q^2 \lambda(\vec{q}_\perp)}{|\vec{u}|} |A^{(\epsilon)}(\vec{q})|^2, \\ \lambda(\vec{q}_\perp) &= \frac{q E_q E_D}{q^2 P^0 - (\vec{q} \cdot \vec{P}) E_q}, \\ E_D &= \sqrt{(\vec{q} - \vec{P})^2 + m_D^2}.\end{aligned}\tag{27}$$

The matrix element $\hat{A}$ is connected with $\hat{S}$ -matrix by relation $\langle f|S|in \rangle = \langle f|in \rangle + i\delta^{(4)}(q_f - q_{in}) \langle f|\hat{A}|in \rangle$.

Now, let us integrate $\frac{d\sigma^\pm}{d\Omega_{\vec{\mu}}}$ (27) over the direction of vector $\vec{\mu}$:

$$\frac{d\sigma^\pm}{d\mu} = \frac{1}{2\pi} \mu \tanh(\pi\mu) \kappa(\mu) \int d\Omega_{\vec{q}} \left\{ \frac{q}{\sqrt{q^2 - q_\perp^2}} P_{-\frac{1}{2}+i\mu} \left(\frac{q}{\sqrt{q^2 - q_\perp^2}} \right) \frac{d\sigma^\pm}{d\Omega_{\vec{q}}} \right\}.\tag{28}$$

Here we used an integral representation of cone function [5]:

$$\int_0^{2\pi} d\varphi (u_0 - \sqrt{u_0^2 - 1} \cos(\varphi - \theta))^{-1/2+i\mu} = 2\pi P_{-1/2+i\mu}(u_0).\tag{29}$$

Subsequent transformation of differential cross-section on $\vec{\mu}$ connect with turning to hyperbolic variables:

$$u = (u_0, u_1, u_2), \quad u_0 = \frac{q}{\sqrt{q^2 - q_\perp^2}}, \quad \vec{u} = \frac{\vec{q}_\perp}{\sqrt{q^2 - q_\perp^2}}, \quad u^2 = u_0^2 - u_1^2 - u_2^2 = 1\tag{30}$$

In this variables the differential volume is:

$$d\Omega_{\vec{q}} = \frac{d\vec{q}_\perp}{q \sqrt{q^2 - q_\perp^2}} = \frac{d\vec{u}}{u_0^3} = \frac{du_0 d\varphi}{u_0^2}\tag{31}$$

where φ is the azimuth angle of vector $\vec{q}_\perp$.

Taking into account that the differential cross-section $\frac{d\sigma^\pm}{d\Omega_{\vec{q}}}$ does not depend on φ we obtain:

$$\frac{d\sigma^\pm}{d\mu} = \mu \tanh(\pi\mu) \kappa(\mu) \int_1^\infty \frac{du_0}{u_0} P_{-\frac{1}{2}+i\mu}(u_0) \left(\frac{d\sigma^\pm}{d\Omega_{\vec{q}}} \right),\tag{32}$$

where the angular part of the cross-section in the right part of integral is expressed through the variable u_0 . Finally we represent the differential cross-section on μ through the differential cross-section on scattering angle. Let θ and φ be axial and azimuth angles of the momentum $\vec{q}$. Let us turn to integrating over this angles in the expression (28). For this we notice that for arbitrary function $f(\vec{q})$ the following integral relation is right:

$$\int f(\vec{q}) d\vec{q} = \int q^2 dq d\Omega f(\vec{q}) = \sum_{\epsilon=\pm 1} \int q^2 dq d\Omega_{\vec{q}} f(\vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}),\tag{33}$$

$$d\Omega = \sin\theta d\theta d\varphi, \quad d\Omega_{\vec{q}} = \frac{1}{q \sqrt{q^2 - q_\perp^2}} d\vec{q}_\perp.$$

From this follows that:

$$\int d\Omega f(\vec{q}) = \sum_{\epsilon=\pm 1} \int d\Omega_{\vec{q}} f(\vec{q}_\perp, \epsilon \sqrt{q^2 - q_\perp^2}),\tag{34}$$

or in terms of θ and φ angles

$$\begin{aligned} \int d\Omega_{\vec{q}} f(\vec{q}_{\perp}, q_3 = +\sqrt{q^2 - q_{\perp}^2}) &= \int_0^1 dz \int_0^{2\pi} d\varphi f(\vec{q}_{\perp}, q_3 = qz) , \\ \int d\Omega_{\vec{q}} f(\vec{q}_{\perp}, q_3 = -\sqrt{q^2 - q_{\perp}^2}) &= \int_{-1}^0 dz \int_0^{2\pi} d\varphi f(\vec{q}_{\perp}, q_3 = qz) , \end{aligned} \quad (35)$$

$$z = \cos \theta , \quad f(\vec{q}_{\perp}, q_3 = qz) \equiv f(\vec{q}) .$$

So relations between variables are

$$qz = \begin{cases} +\sqrt{q^2 - q_{\perp}^2} & \text{forward half-sphere,} \\ -\sqrt{q^2 - q_{\perp}^2} & \text{backward half-sphere.} \end{cases} , \quad q_{\perp} = q\sqrt{1 - z^2} .$$

We turn to integrating over angles in the integral (28). We obtain:

$$\begin{aligned} \frac{d\sigma^+}{d\mu} &= \frac{1}{2\pi} \mu \tanh(\pi\mu) \kappa(\mu) \int_0^1 \frac{dz}{z} P_{-1/2+i\mu}\left(\frac{1}{z}\right) \frac{d\sigma}{dz} , \\ \frac{d\sigma^-}{d\mu} &= \frac{1}{2\pi} \mu \tanh(\pi\mu) \kappa(\mu) \int_{-1}^0 \frac{dz}{|z|} P_{-1/2+i\mu}\left(\frac{1}{|z|}\right) \frac{d\sigma}{dz} . \end{aligned} \quad (36)$$

where $\frac{d\sigma}{dz}$ is the differential cross-section on the cosine of scattering angle of particle C in the $A + B \rightarrow C + D$ process. Here is taken into account that $\frac{d\sigma^{\pm}}{d\Omega_{\vec{q}}}$ do not depend of variable φ , therefore

$$\frac{d\sigma^{\pm}}{d\Omega_{\vec{q}}} = \frac{1}{2\pi} \frac{d\sigma}{dz} .$$

Unifying relations (22) , (27) , (36) and taking into account that:

$$\frac{1}{(2\pi)^2} \int \kappa(\mu) \bar{\xi}(\vec{q}_{\perp}, \vec{\mu}) d\Omega_{\vec{\mu}} = 1 ,$$

we obtain the norm of the differential cross-section on the total cross-section $\sigma(AB \rightarrow CD)$

$$\sum_{\epsilon=\pm 1} \int_0^{\infty} \left(\frac{d\sigma^{(\epsilon)}}{d\mu} \right) d\mu = \int_{-1}^1 \left(\frac{d\sigma}{dz} \right) dz = \frac{N}{TVn_1n_2|\vec{u}|} = \sigma(AB \rightarrow CD) . \quad (37)$$

As follows from (27), in contrast to $\frac{d\sigma^{\pm}}{d\Omega_{\vec{q}}}$ the differential cross-section $\frac{d\sigma^{\pm}}{d\Omega_{\vec{\mu}}}$ is not positively sign determined on all μ interval. Contribution of negative value area of $\frac{d\sigma^{\pm}}{d\Omega_{\vec{\mu}}}$ reduces effectively to decreasing of the total event number of particle C creation. So, this spatial area we can interpret as area where taking place an absorption of particles C . But the total number of asymptotic states with define μ regulates by the relation (37).

APPLICATION TO SIMPLE MODELS $\frac{d\sigma}{d\Omega}$ IN C.M.F.

As an example we discuss an one-particle exchange in t -channel, for elastic $A+B \rightarrow A+B$ scattering. Corresponding cross section as function of u_0 has a polar view:

$$\frac{d\sigma}{d\Omega} = N_0 \frac{\alpha^2}{(t - M^2)^2} = \frac{\alpha^2 N_0}{(2q^2)^2} \frac{1}{(z_0 - z)^2} = \frac{\alpha^2 N_0}{(2z_0 q^2)^2} \frac{u_0^2}{(u_0 - \frac{\epsilon}{z_0})^2} , \quad (38)$$




FIG. 1: a) Distribution function on b in the model of one particle exchange. $bq < 1/2$ – area forbidden by uncertainty relation, $1/2 < bq \lesssim \sqrt{2}$ – area of creation particles C , $bq \gtrsim \sqrt{2}$ – area of absorption particles C . b) The zone structure of plain $\vec{b}$ in the model of one particle exchange. Zone I – area forbidden by uncertainty relation, zone II – area of creation particles C , zone III – area of absorption particles C .

$$z_0 = 1.001$$

here

$$z_0 = 1 + \frac{2M^2 s}{\lambda(s, m_A^2, m_B^2)} = 1 + \frac{M^2}{2q^2},$$

M – mass of exchange particle, $\lambda(x, y, z) = (x^2 + y^2 + z^2 - 2xy - 2xy - 2yz)$ – wellknown the triangle function, α – the coupling constant,

$$N_0 = (2\pi)^2 \frac{q^2 \tilde{\lambda}(\vec{q}_\perp)}{|\vec{u}|} = (2\pi)^2 \frac{(s^2 - (m_1^2 - m_2^2)^2)^2}{16s^3}.$$

In this case for the cross-section $\sigma^\pm$ we obtain:

$$\sigma^\pm = \frac{\alpha^2 N_0}{(2q^2)^2} \frac{2\pi}{z_0(z_0 \mp 1)}, \quad (39)$$

where

$$\sigma^{(\varepsilon)} = \int_0^\infty \left(\frac{d\sigma^{(\varepsilon)}}{d\mu} \right) d\mu = \begin{cases} \int_0^1 \left(\frac{d\sigma}{dz} \right) dz, & \varepsilon = 1 \\ \int_{-1}^0 \left(\frac{d\sigma}{dz} \right) dz, & \varepsilon = -1 \end{cases}. \quad (40)$$

The integral (32) with such cross section is computed analytically. The normalize distribution on nearest approach parameter takes a view:

$$\frac{1}{\sigma^\varepsilon} \frac{d\sigma^\varepsilon}{db} = \frac{q^2 b \tanh(\pi\mu) \kappa(\mu)}{2 \cosh(\mu\pi)} \frac{z_0 - \varepsilon}{z_0} \left[P_{i\mu-1/2} \left(\frac{-\varepsilon}{z_0} \right) + \frac{\varepsilon}{\sqrt{z_0^2 - 1}} P_{i\mu-1/2} \left(\frac{-\varepsilon}{z_0} \right) \right], \quad (41)$$

where $\varepsilon = \pm 1$, and $P_{i\mu-1/2}^1(x)$ – associated cones function. The plot of (41) is shown on Fig.1a.

Since $z_0 \geq 1$ so the argument of cones functions belong to segment $[-1, 1]$. The cones function is positive with such argument (and associated cones functions with $m > 0$ too). So the cross-section is positive defined on $\varepsilon = 1$. On $\varepsilon = -1$ the $\vec{b}$ -plane divides into zones with radiuses R_0 , R_1 and R_2 Fig.1b.

Here $R_0^2 = \hbar^2/4q^2$ defines a border of forbidden area, where Geizenberg uncertainty relation is broken (phase space of particle C is less then allowable). Area $R_0 < b < R_1$ defines spatial area of particle C creation. And area $b > R_1$ is area where absorption of particles is taken place relating to equality (37). At that, if we symbolically take density of distribution $I(b) \sim 1$ in area $R_0 < b < R_1$, when density in area $b > R_1$ would be $\sim 10^{-5}$. It is made conditional

upon that exponential decrease of $I(b)$ in the backward sphere determines by radius $R \sim 1/q$ i.e. $I(b) \sim \exp(-\pi bq)$ at $bq \gg 1$.

The left border of R_2 -zone is defined by zero of expression (41). In the area of small values of parameter $\frac{M^2}{s}$ the cross-section (41) turn into zero on

$$\mu = \mu_0 = \frac{\sqrt{7}}{2} + 2\frac{2}{\sqrt{7}}\epsilon + \frac{2}{21\sqrt{7}}\epsilon^2 + O(\epsilon^3), \quad \epsilon = \frac{2M^2 s}{\lambda(s, m_1^2, m_2^2)} \sim \frac{M^2}{s}.$$

In the area $\frac{M^2}{s} \ll 1$ the cross-section takes a view:

$$\frac{1}{\sigma^-} \frac{d\sigma^-}{db} = q^2 b \frac{\tanh(\pi\mu)\kappa(\mu)}{\cosh(\mu\pi)} \left[\frac{7-4\mu^2}{8} + O\left(\frac{M^2}{s}\right) \right], \quad (42)$$

$$\frac{1}{\sigma^+} \frac{d\sigma^+}{db} = q^2 b \frac{\tanh(\pi\mu)\kappa(\mu)}{\cosh(\mu\pi)} \left[\frac{\cosh(\mu\pi)}{\pi} + O\left(\frac{M^2}{s}\right) \right]. \quad (43)$$

CONNECTION BETWEEN $\langle b^2 \rangle$ AND THE CROSS-SECTION ON TRANSVERSE MOMENTUM OF PARTICLE C IN C.M.F OF A AND B PARTICLES

Let us calculate the an average value of nearest approach parameter square $\langle b_{\pm}^2 \rangle$. We have by definition:

$$\langle \mu_{\pm}^2 \rangle = \frac{1}{\sigma^{\pm}} \int_0^{\infty} \mu^2 \frac{d\sigma^{\pm}}{d\mu} d\mu, \quad (44)$$

where $\sigma^{\pm}$ was defined in (40). So:

$$\langle b_{\pm}^2 \rangle = \frac{1}{q^2} \left(\langle \mu_{\pm}^2 \rangle + \frac{1}{4} \right), \quad q = \tilde{q}. \quad (45)$$

Substituting the representation for $\frac{d\sigma^{\pm}}{d\mu}$ from (32), into (44) and taking into account that (compare with (22) and (29)):

$$\kappa(\mu) = 2\pi \int_1^{\infty} P_{i\mu-1/2}(x) \frac{dx}{x},$$

we obtain

$$\langle b_{\pm}^2 \rangle = \frac{2\pi}{\tilde{q}^2 \sigma^{\pm}} \int (\mu^2 + \frac{1}{4}) P_{i\mu-1/2}(x) P_{i\mu-1/2}(u_0) \frac{d\sigma^{\pm}}{d\Omega} \frac{dx}{x} \frac{du_0}{u_0} d\Omega_{\mu}. \quad (46)$$

The differential equation on cones functions $P_{i\mu-1/2}(u_0)$ with argument u_0 can be represented in following form [5]:

$$(\mu^2 + \frac{1}{4}) P_{i\mu-1/2}(u_0) = \left(-(u_0^2 - 1) \frac{d^2}{du_0^2} - 2u_0 \frac{d}{du_0} \right) P_{i\mu-1/2}(u_0).$$

The left part of this expression went as underintegral factor into relation (46) and its substitution allows us integrate over μ . From the completeness relation of cones functions follow [2]:

$$\int_0^{\infty} P_{i\mu-1/2}(x) P_{i\mu-1/2}(u_0) d\Omega_{\mu} = \delta(x - u_0). \quad (47)$$

Next, we can integrate over variable x . After some transformations we obtain:

$$\langle b_{\pm}^2 \rangle = \frac{2\pi}{\tilde{q}^2 \sigma^{\pm}} \int \frac{2}{u_0^3} \frac{d\sigma^{\pm}}{d\Omega} \frac{du_0}{u_0}. \quad (48)$$

Crossing in this relation from the variable u_0 to the scattering angle θ , we obtain:

$$\langle b_\varepsilon^2 \rangle = \begin{cases} \frac{2}{q^2 \sigma^+} \int_0^1 \cos^2 \theta \frac{d\sigma^+}{d \cos \theta} d \cos \theta, & \varepsilon = 1 \\ \frac{2}{q^2 \sigma^-} \int_{-1}^0 \cos^2 \theta \frac{d\sigma^-}{d \cos \theta} d \cos \theta, & \varepsilon = -1 \end{cases}. \quad (49)$$

So, finally we have:

$$\langle b_\pm^2 \rangle = \frac{8s}{\lambda(s, m_C^2, m_D^2)} \langle \cos^2 \theta_\pm \rangle. \quad (50)$$

where

$$\langle \cos^2 \theta_\varepsilon \rangle = \begin{cases} \frac{1}{\sigma^+} \int_0^1 z^2 \frac{d\sigma}{dz} dz, & \varepsilon = 1 \\ \frac{1}{\sigma^-} \int_{-1}^0 z^2 \frac{d\sigma}{dz} dz, & \varepsilon = -1 \end{cases}, \quad z = \cos \theta,$$

and $\langle b_\pm^2 \rangle$ is defined by relation (44).

It follows:

$$\langle \mu_\pm^2 \rangle = 2 \langle \cos^2 \theta_\pm \rangle - \frac{1}{4}. \quad (51)$$

The parameter μ is real number, so $\mu^2 > 0$. From it and (51) follows an important physical inequality:

$$\langle \cos^2 \theta_\pm \rangle \geq \frac{1}{8}. \quad (52)$$

Let us discuss the nature of this inequality. It follows from the reality of parameter $\mu = \sqrt{b^2 q^2 / \hbar^2 - 1/4}$, where $\hbar$ – Plank constant. Here b^2 is an eigen value of Kazimir's operator on $SO(2, 1)$ -group [2]. Spectrum of this operator in Hilbert space of states satisfies condition $b^2 q^2 \geq \hbar^2/4$, and that is provide a reality of parameter μ . By itself this inequality has an quantum nature and show us the fact that particle C can not creates in phase space less when it allows by Geizenberg's uncertainty relation.

We notice on the fact that relation (50) between $\langle b_\pm^2 \rangle$ and $\langle \cos^2 \theta_\pm \rangle$ right for any process $A + B \rightarrow C + D$ in center of mass frame A and B particles.

In the model with one-particle exchange (38) it is easy to obtain that

$$\langle \cos^2 \theta_\pm \rangle = \pm 2z_0^2(z_0 \mp 1) \ln \left(\frac{z_0 \mp 1}{z_0} \right) \mp z_0 + 2z_0^2.$$

So it follows:

$$\langle b_\pm^2 \rangle = \frac{1}{q^2} \left[\pm 4z_0^2(z_0 \mp 1) \ln \left(\frac{z_0 \mp 1}{z_0} \right) \mp 2z_0 + 4z_0^2 \right].$$

Let us analyze this expressions as functions of parameter $z_0 = 1 + M^2/2q^2$ (relation (38)). The analyze shows that $\langle \cos^2 \theta_\pm \rangle$ changes in borders:

$$\frac{1}{3} < \langle \cos^2 \theta_+ \rangle \leq 1,$$

$$0.23 \lesssim \langle \cos^2 \theta_- \rangle < \frac{1}{3},$$

when z_0 changes in interval from 1 to ∞ .

DIFFERENTIAL CROSS SECTION ON NEAREST APPROACH PARAMETER $\vec{b}$ IN L.F.

As it was note in the previous section, in the labor frame of B particle energy and momentum of C particle depend of scattering angle (13). Consequence of this fact is appearance on the reaction plane $q = \tilde{q}$ as the parameter μ dependence from the scattering angle, if we use relation (26) for the transition to the parameter b . $SO_\mu(2, 1)$ algebra implements on basis functions, for which the parameter μ is natural variable. Completeness and orthogonality of basis are appeared in terms of it. If we take μ in relation (24) as an independent parameter, then differential cross section $\frac{d\sigma}{d\mu}$ is defined unambiguously and doesn't depend of frame. But at the same time the physical meaning of this parameter isn't clear, only as phase space measure $(b, q_\perp)$. In c.m.f. transition in the integral (24) from μ to b does not change anything considerably, and differential distributions on μ and on b are equivalent. But in l.f. the differential distribution changes radically. Let us cross in the integral (24) to the variable $\vec{b}$ using relations (26). Than we have:

$$N = \frac{1}{(2\pi)^2} \text{Re} \sum_{\epsilon=\pm 1} \int d\vec{b} \int d\Omega_{\vec{q}} \vartheta[b - R_0(\tilde{q})] q^2 \tanh(\pi\mu) \kappa(\mu) \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \frac{dN^{(\epsilon)}}{d\Omega_{\vec{q}}} , \quad (53)$$

here

$$\mu = (b^2 q^2 - \frac{1}{4})^{1/2} , \quad R_0(\tilde{q}) = \frac{\hbar}{2\tilde{q}} , \quad \vartheta(x) - \text{discontinuous function} , \quad q = \tilde{q} ,$$

and $\tilde{q} = p_C$ defines in (13)

From this it follows for the differential cross section on $\vec{b}$:

$$\frac{d\sigma^{(\epsilon)}}{d\vec{b}} = \frac{1}{(2\pi)^2} \text{Re} \sum_{\epsilon=\pm 1} \int d\Omega_{\vec{q}} \vartheta[b - R_0(\tilde{q})] q^2 \tanh(\pi\mu) \kappa(\mu) \bar{\xi}(\vec{q}_\perp, \vec{\mu}) \frac{d\sigma^{(\epsilon)}}{d\Omega_{\vec{q}}} . \quad (54)$$

Integrating left and right part over the direction of $\vec{b}$ vector and crossing to angle variables θ and φ of $\vec{q}_\perp$ vector we obtain the result, which is analogous to relations (36):

$$\begin{aligned} \frac{d\sigma^+}{db} &= \frac{b}{2\pi} \int_0^1 \frac{dz}{z} \vartheta[b - R_0(\tilde{q})] q^2 \tanh(\pi\mu) \kappa(\mu) P_{-1/2+i\mu}\left(\frac{1}{z}\right) \frac{d\sigma}{dz} , \\ \frac{d\sigma^-}{db} &= \frac{b}{2\pi} \int_{-1}^0 \frac{dz}{|z|} \vartheta[b - R_0(\tilde{q})] q^2 \tanh(\pi\mu) \kappa(\mu) P_{-1/2+i\mu}\left(\frac{1}{|z|}\right) \frac{d\sigma}{dz} . \end{aligned} \quad (55)$$

From comparison of (55) and (36) it follows that differences between distributions on the parameter b in c.m.f. and in l.f. generates by difference of the dependence of particle C momentum from kinematic variables s and t for this two cases. From the explicit expression $\tilde{q} = p_C$ (36) it follows that it is a fluent function of angle and with certain accuracy it is possible cross from $\tilde{q}$ to some midvalue in integrals (55). For example, we receive that on the backward semisphere:

$$\tilde{q}(z) \simeq \tilde{q}(\bar{z}) , \quad z = \cos(\theta) , \quad \bar{z} = -0.5 ,$$

and it isn't depend on angle. Then the distribution (55) coincides with the distribution (36) at $\tilde{q}(z) \simeq \tilde{q}(\bar{z})$. More detailed structure of particle C creation area reflects in a generalized distribution function:

$$\frac{d^2\sigma}{db dz} = \frac{b}{2\pi} \cdot \vartheta[b - R_0(\tilde{q})] q^2 \tanh(\pi\mu) \kappa(\mu) \frac{1}{|z|} P_{-1/2+i\mu}\left(\frac{1}{|z|}\right) \frac{d\sigma}{dz} , \quad -1 < z < 1 , \quad (56)$$

$$q = \tilde{q} = p_C(z) , \quad \mu = (b^2 q^2 - \frac{1}{4})^{1/2} .$$

It gives us a tomographic picture of partial integrals over z to the spatial distribution $\frac{d\sigma}{db}$ in the whole interval of b . It is similar to Wigner function [6] From it we can get expressions for the differential cross section on $q_\perp$ (11), for differential cross section on b (36, 55) and norm relations (37).




FIG. 2: a) Differential cross section of process $\gamma + p \rightarrow \pi^0 + p$, $x = \cos \theta$
b) Differential cross section on b for process $\gamma + p \rightarrow \pi^0 + p$




FIG. 3: Generalized distribution function in plane (b, z) for the process $\gamma + p \rightarrow \pi^0 + p$, $E_\gamma = 5 \text{ GeV}$, $q \approx 665 \text{ MeV}$

As an illustration of represented method, let us consider process of neutral π -meson photoproduction on proton. There is experimental data on angle distribution of π^0 in the interval $-0.75 < z = \cos \theta < 0$ (backward semisphere) at photon energy $E_\gamma = 5 \text{ GeV}$ (Fig.3a)[7]. At such energy and in this angle interval the meson energy depends of angles weakly. And we receive in relations (56) π^0 momentum as its midvalue $q \approx 665 \text{ MeV}$. There is the generalized distribution function at current parameters (Fig.4). The area I is defined by condition $b < R_0 = \hbar/2q$. This is lower border of parameter b physical values, it is consequent of Geizenberg uncertainty relation in phase space (b, q) . $R_0 = 1.498 \cdot 10^{-14} \text{ cm}$ at current values. The area II is particle C creation area. In this area the distribution function is positive. Finally, the area III is area where the distribution function is negative. Contribution of this area to the total number of particle C creation events is negative. So, as it was discussed, this area is interpreted as the particle C absorption area. The total number of events in unit time is regulate according to (37).

Fig.3b is distribution on b integrated over available angle interval $-0.75 < z = \cos \theta < 0$:

$$\frac{d\sigma}{db} = \int_{-0.75}^0 \left[\frac{d^2\sigma}{db dz} \right] dz .$$

Evidently to the graphic, particle C creation areas are also divide to zones. Numerical calculation at current parameters gives us the following values of zone radiuses. The first zone - ($R_0 < b < R_1$), where $R_1 = 5.982 \cdot 10^{-14} \text{ cm}$. The second zone - ($R_2 < b < R_3$), where $R_2 = 1.757 \cdot 10^{-13} \text{ cm}$, $R_3 = 2.79 \cdot 10^{-13} \text{ cm}$. The third zone - ($b > R_4$), where $R_4 = 3.899 \cdot 10^{-13} \text{ cm}$ and so on.

The cross section fast falls with b growing. If we take the maximum value of cross section in the first zone as unit, when the maximum value in the first absorption zone ($bq \sim 4$) would be equal to $1.5 \cdot 10^{-3}$, and the maximum in the

second creation zone would be $2.8 \cdot 10^{-6}$. Here we represented estimates of the spatial distribution starting from some midvalue of π^0 energy. In fact we have to do exact calculation in (56) of μ angle dependence, using corresponding relation for π^0 momentum.

CONCLUSION

So, in the paper it was relieved the exact relation between the differential cross section on the detect particle momentum and the distribution function on out going nearest approach parameter of this particle. This function describes a distribution of matter in reaction area and allows understand the nature of its composite parts on hadronic scale. This connection has a different view in the center of mass frame of A and B particles and in laboratory frame of particle B . On the simple model it was shown that reaction area divides on creation and absorption areas of particle C . It was received a relation between $\langle b_{\pm}^2 \rangle$ and $\langle \cos^2 \theta_{\pm} \rangle$ in center of particles A and B mass frame, where average is going on corresponding differential cross-sections. This relation is exact and right for any $A + B \rightarrow C + D$ processes. There is a bottom bound on $\langle \cos^2 \theta_{\pm} \rangle$ also follows from it.

Only completeness of states $|\vec{\mu}, q, \epsilon \rangle$ in one particle Fock space was used for finding connection between cross sections.

It is received the generalized distribution on plane (b, z) , and the differential cross sections and the norm follows from that. As an illustration it was considered a real process of photoproduction π^0 on the proton, and it was done valuation of the main characteristics of spatial distribution.

ACKNOWLEDGEMENTS

It is our pleasure to thank prof. I.B.Khriplovich for constructive discussion and critical remarks. This investigation has been supported in part by grant President of Russian Federation for support of leading scientific schools (NSH -5362.2006.2) (N.B & A.N.V), by the Russian Foundation for Fundamental Research (A.A.V), and by the Heisenberg-Landau Program of JINR grant No. 07-02-91557 (A.A.V)

* Electronic address: vall@irk.ru

† Electronic address: Maxim.Polyakov@tp2.ruhr-uni-bochum.de

‡ Electronic address: avlad@theor.jinr.ru

- [1] N.N.Bogolubov, D.V.Shirkov, *Introduction to the Theory of Quantized Fields* Chichester: Willey-InterScience, 3rd ed. 1980
- [2] M.V.Polyakov, O.N.Soldatenko, A.N.Vall, A.A.Vladimirov, arXiv:0708.2857v1 [hep-ph]
- [3] V.I. Gol'danskii, I.L. Rozental, Yu P. Nikitin, *Kinematic Methods in High Energy Physics*, Harwood Academic (Medical, Reference and Social Sc), 1989
- [4] I. S. Gradshtein and I. M. Ryzhik, *Tables of Integrals, Series, and Functions*, Pergamon, New York, 1964.
- [5] Bateman H. and Erdelyi A., *Higher Transcendental Functions* vol.1 McGraw-Hill, New York (1953).
- [6] E.P. Wigner, Phys.Rev.40 (1932) 749.
P.Carruthers, F.Zachariasen, Rev.Mod.Phys.55, N1, (1983), 283
- [7] R.L.Anderson *et al.* Phys.Rev.D14:679,1976.